



SEER Incidence - AA Rates for Additional Races/Registries, 1992-2001

 [Back to referring page](#)

Selections:

Statistic type = Age-adjusted rate;
Standard population = 2000 U.S.;
SEER registry = Total (registries depend on race/ethnicity);
Site = Lung and Bronchus;
Race/ethnicity = All races;
Age at diagnosis = All ages;

Results: [D]



Notes:

http://canques.seer.cancer.gov/cgi-bin/cq_submit?dir=seer2001&db=2&rpt=LINE&sel=1^... 7/12/2004

PM3003723889

- Statistics were generated from malignant cases only.
- Rates are expressed as cases per 100,000.
- Statistics are provided by the SEER Program for research purposes only.
- Hispanic and Non-Hispanic are not mutually exclusive from White, Black, American Indian/Alaska Native, and Asian or Pacific Islander.
- Statistics for Hispanics and Non-Hispanics do not include cases from the Detroit, Hawaii, and Alaska Natives registries.

Suggested Citation - Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Databases: Incidence - SEER 11 Regs + AK Public-Use, Nov 2003 Sub for Expanded Races (1992-2001) and Incidence - SEER 11 Regs Public-Use, Nov 2003 Sub for Hispanics (1992-2001), National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released April 2004, based on the November 2003 submission.

Line Chart Variables:

X-Axis = Year of diagnosis;
Series = Sex;

Alternative Formats:

[Vertical Bar Chart](#) | [Table](#) | [Delimited Data](#)



[Back to referring page](#)

Software provided by SEER: cq_submit 3.1

Comments to: seerweb@imsweb.com